

DTaP (Diphtheria, Tetanus, Pertussis) Vaccine: *What You Need to Know*

Many Vaccine Information Statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1 Why get vaccinated?

DTaP vaccine can prevent **diphtheria, tetanus, and pertussis**.

Diphtheria and pertussis spread from person to person. Tetanus enters the body through cuts or wounds.

- **DIPHtheria (D)** can lead to difficulty breathing, heart failure, paralysis, or death.
- **TETANUS (T)** causes painful stiffening of the muscles. Tetanus can lead to serious health problems, including being unable to open the mouth, having trouble swallowing and breathing, or death.
- **PERTUSSIS (aP)**, also known as “whooping cough,” can cause uncontrollable, violent coughing which makes it hard to breathe, eat, or drink. Pertussis can be extremely serious in babies and young children, causing pneumonia, convulsions, brain damage, or death. In teens and adults, it can cause weight loss, loss of bladder control, passing out, and rib fractures from severe coughing.

2 DTaP vaccine

DTaP is only for children younger than 7 years old. Different vaccines against tetanus, diphtheria, and pertussis (Tdap and Td) are available for older children, adolescents, and adults.

It is recommended that children receive 5 doses of DTaP, usually at the following ages:

- 2 months
- 4 months
- 6 months
- 15–18 months
- 4–6 years

DTaP may be given as a stand-alone vaccine, or as part of a combination vaccine (a type of vaccine that combines more than one vaccine together into one shot).

DTaP may be given at the same time as other vaccines.

3 Talk with your health care provider

Tell your vaccine provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of any vaccine that protects against tetanus, diphtheria, or pertussis**, or has any **severe, life-threatening allergies**.
- Has had a **coma, decreased level of consciousness, or prolonged seizures within 7 days after a previous dose of any pertussis vaccine (DTP or DTaP)**.
- Has **seizures or another nervous system problem**.
- Has ever had **Guillain-Barré Syndrome** (also called GBS).
- Has had **severe pain or swelling after a previous dose of any vaccine that protects against tetanus or diphtheria**.

In some cases, your child's health care provider may decide to postpone DTaP vaccination to a future visit.

Children with minor illnesses, such as a cold, may be vaccinated. Children who are moderately or severely ill should usually wait until they recover before getting DTaP.

Your child's health care provider can give you more information.



U.S. Department of
Health and Human Services
Centers for Disease
Control and Prevention

4 Risks of a vaccine reaction

- Soreness or swelling where the shot was given, fever, fussiness, feeling tired, loss of appetite, and vomiting sometimes happen after DTaP vaccination.
- More serious reactions, such as seizures, non-stop crying for 3 hours or more, or high fever (over 105°F) after DTaP vaccination happen much less often. Rarely, the vaccine is followed by swelling of the entire arm or leg, especially in older children when they receive their fourth or fifth dose.
- Very rarely, long-term seizures, coma, lowered consciousness, or permanent brain damage may happen after DTaP vaccination.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

5 What if there is a serious problem?

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at www.vaers.hhs.gov or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff do not give medical advice.*

6 The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Visit the VICP website at www.hrsa.gov/vaccinecompensation or call **1-800-338-2382** to learn about the program and about filing a claim. There is a time limit to file a claim for compensation.

7 How can I learn more?

- Ask your health care provider.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636** (**1-800-CDC-INFO**) or
 - Visit CDC's website at www.cdc.gov/vaccines

Vaccine Information Statement (Interim)
DTaP (Diphtheria, Tetanus,
Pertussis) Vaccine



Office use only

04/01/2020 | 42 U.S.C. § 300aa-26